



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 1990

---

## **Improved cardiac performance with human calcitonin gene related peptide in patients with congestive heart failure**

Gennari, C ; Nami, R ; Agnusdei, D ; Fischer, J A

**Abstract:** Study objective - The aim of the study was to assess the cardiovascular effects of human calcitonin gene related peptide (CGRP) in patients with congestive heart failure. Design - The effects of CGRP II (or ),  $12.5 \text{ g} \cdot \text{h}^{-1}$ , given by intravenous infusion for 24 h to digitalised patients with congestive heart failure, were assessed by measurement of cardiac functional indices. Patients - Five patients (four female) were studied. Age was 73-82 years. Three were in New York Heart Association phase III and two in phase IV. Measurements and main results - The pre-ejection period to left ventricular ejection time ratio and the QT distance adjusted for heart rate were lowered by 21% and 4% respectively. The left ventricular shortening index was raised by 43%. The arterial pressure and heart rate did not change consistently. Conclusion - Calcitonin gene related peptide improves myocardial contractility in patients with congestive heart failure. This is the first time this has been shown

DOI: <https://doi.org/10.1093/cvr/24.3.239>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-154807>

Journal Article

Published Version

Originally published at:

Gennari, C; Nami, R; Agnusdei, D; Fischer, J A (1990). Improved cardiac performance with human calcitonin gene related peptide in patients with congestive heart failure. *Cardiovascular Research*, 24(3):239-241.

DOI: <https://doi.org/10.1093/cvr/24.3.239>

# Improved cardiac performance with human calcitonin gene related peptide in patients with congestive heart failure

C Gennari, R Nami, D Agnusdei, J A Fischer

## Abstract

**Study objective** – The aim of the study was to assess the cardiovascular effects of human calcitonin gene related peptide (CGRP) in patients with congestive heart failure.

**Design** – The effects of CGRP II (or  $\beta$ ), 12.5  $\mu\text{g}\cdot\text{h}^{-1}$ , given by intravenous infusion for 24 h to digitalised patients with congestive heart failure, were assessed by measurement of cardiac functional indices.

**Patients** – Five patients (four female) were studied. Age was 73–82 years. Three were in New York Heart Association phase III and two in phase IV.

**Measurements and main results** – The pre-ejection period to left ventricular ejection time ratio and the QT distance adjusted for heart rate were lowered by 21% and 4% respectively. The left ventricular shortening index was raised by 43%. The arterial pressure and heart rate did not change consistently.

**Conclusion** – Calcitonin gene related peptide improves myocardial contractility in patients with congestive heart failure. This is the first time this has been shown.

Congestive heart failure is usually caused by reduced cardiac output as a result of impaired myocardial contractility. Its treatment includes positive inotropic agents, eg, digitalis, as well as vasodilator and diuretic therapy.<sup>1</sup> Calcitonin gene related peptide (CGRP) is a recently discovered neuropeptide with potent cardiovascular effects that include positive chronotropic and inotropic actions on the heart,

vasodilatation, and hypotension.<sup>2–9</sup> The stimulation of the heart rate and coronary vasodilatation presumably depend on the local release of CGRP from afferent nerve fibres in the heart induced by capsaicin and interaction with receptors linked to cyclic adenosine monophosphate (AMP).<sup>10–16</sup> The improved ventricular contractility of CGRP was suppressed in normal human subjects through blockade of adrenergic receptors, and may be driven by reflex sympathetic stimulation.<sup>4</sup> Here we show evidence of improved ventricular contractility with CGRP as recorded with non-invasive techniques in five digitalised patients with congestive heart failure.

## Methods

We studied four female and one male patient aged 73 to 82 years with congestive heart failure, three in New York Heart Association phase III and two in phase IV.<sup>17</sup> Four patients had coronary artery disease and the fifth had hypertrophic cardiomyopathy. Body weight was 44–66 kg, and height 147–158 cm. Informed consent about the experimental treatment with CGRP was obtained from all patients, and the study was approved by the appropriate ethics committee.

The patients were treated with oral  $\beta$ -methyl digoxin (Lanitop, Boehringer, Mannheim, W Germany), 0.6 mg (0.77 nmol) per day for three days, followed by 0.3 mg (0.39 nmol) per day for four days (figure), after which they were given human CGRP II (or  $\beta$ ) (Peninsula Laboratories, Belmont, CA, USA), 12.5  $\mu\text{g}\cdot\text{h}^{-1}$  (3.3 nmol) by intravenous minipump infusion for 24 h.

The systemic arterial pressure was monitored continuously by intra-arterial manometry. The duration of the pre-ejection period and the left ventricular ejection time were determined using electrocardiography, phonocardiography, and external carotid pulse tracings.<sup>18</sup> The corrected QT interval ( $\text{QT}_c$ ) was calculated by dividing QT by the square root of the preceding R-R interval. The left ventricular cavity was measured before and during the infusions with CGRP by M mode echocardiography at the end diastolic and the end systolic filling times.<sup>19</sup> The left ventricular shortening index was calculated as the ratio between the left ventricular end diastolic minus the end systolic diameters, and the left ventricular end diastolic diameter.

Institute of Medical Semeiotics, University of Siena, 53100 Siena, Italy

C Gennari

R Nami

D Agnusdei

Research Laboratory for Calcium Metabolism, Departments of Orthopaedic Surgery and Medicine, University of Zurich, Forchstrasse 340, 8008 Zurich, Switzerland.

J A Fischer

Correspondence to: Professor Fischer

**Key words:** calcitonin gene related peptide; contractility; heart failure

Submitted 10 May 1989

Accepted 31 October 1989

Serum concentrations of digoxin were measured by radioimmunoassay.<sup>20</sup> Total serum calcium was measured by atomic absorption spectrophotometry (Perkin-Elmer, Norwalk, CN, USA).<sup>21</sup>

Statistical analysis was by paired *t* test.<sup>22</sup> Values are given as means (SEM).

## Results

Congestive heart failure was treated with  $\beta$ -methyl digoxin and therapeutic serum concen-


trations ( $0.7\text{--}1.8\text{ nmol}\cdot\text{litre}^{-1}$ ) were achieved (figure).<sup>20</sup> The heart rate was lowered ( $p<0.05$ ), but the pre-ejection period to left ventricular ejection time ratio and the  $QT_c$  intervals were only minimally decreased, from  $0.50$  (SEM  $0.02$ ) to  $0.47(0.02)$  ( $p<0.01$ ), and from  $0.43(0.03)$  s to  $0.42(0.03)$  s, respectively ( $p<0.01$ ). Myocardial contractility remained largely unchanged during the seven day treatment with therapeutic doses of  $\beta$ -methyl digoxin. As a result, the patients were essentially refractory to treatment with digoxin. No other treatment was used.

Human CGRP II was then given as an intravenous infusion for 24 h at a rate previously used in normal subjects.<sup>23</sup> The arterial pressure was minimally lowered and the heart rate remained slightly raised ( $p>0.05$ ), but the pre-ejection period to left ventricular ejection time ratio was lowered from  $0.47(0.03)$  to  $0.37(0.03)$  (normal  $<0.35$ ;  $p<0.001$ ), and the  $QT_c$  intervals were shortened from  $0.42(0.03)$  to  $0.41(0.03)$  s ( $p<0.01$ ). The left ventricular shortening index was raised from  $0.28(0.02)$  to  $0.42(0.04)$  (normal  $>0.29$ ;  $p<0.01$ ). Improved contractility of the heart ventricle was evident with CGRP in all the five patients. With the dose of CGRP used, the undesired hypotensive and positive chronotropic effects were limited and did not affect the therapeutic efficacy of CGRP on ventricular contractility. All five patients noted subjective improvements. With CGRP, serum calcium levels remained unchanged: before CGRP  $2.29(0.03)$  mmol·litre<sup>-1</sup>; after CGRP  $2.29(0.04)$  mmol·litre<sup>-1</sup>.

## Discussion

The improvement of myocardial contractility is a principal object of the treatment of congestive heart failure. Digitalis has limited efficacy and potential toxicity.  $\beta$  Adrenergic agonists are thought to exert positive inotropic effects on the heart through stimulation of cyclic AMP production.<sup>1</sup> As a result, calcium influx is stimulated in myocardial cells.<sup>24</sup> The improved ventricular contractility of CGRP was suppressed in normal subjects through blockade of adrenergic receptors, and may result in part from reflex sympathetic stimulation.<sup>4</sup> Coronary vasodilatation is achieved through interaction of CGRP with its receptors and stimulation of cyclic AMP production.<sup>6, 7, 13, 15, 16</sup>

Here we have used non-invasive procedures. Reduction of systolic time intervals is brought about through positive inotropic agents and reduction in afterload, eg, due to vasodilatation.<sup>18</sup> The combined positive inotropic action on the heart and vasodilator properties of CGRP explain the improvement of congestive heart failure in the present patients. In view of the vasodilator activity of CGRP it remains to be shown whether cardiac function is also ameliorated through the positive inotropic action of the peptide<sup>25</sup> in patients with congestive heart failure.



Effects of peroral (po)  $\beta$ -methyl digoxin and intravenous (iv) human CGRP II (hCGRP II) on arterial pressure, heart rate, pre-ejection period to left ventricular ejection time ratio (PEP/LVET), the  $QT_c$  distance adjusted for heart frequency ( $QT_c$ ), the left ventricular shortening index (LVSI), and serum concentrations of digoxin in five patients with congestive heart failure. Each value represents the mean, bars = SEM. Open symbols represent statistically significant changes from the values obtained at the start of the study, or in the case of the left ventricular shortening index from before the administration of CGRP ( $\circ$ ,  $p<0.05$ ;  $\triangle$ ,  $p<0.01$ ;  $\square$ ,  $p<0.001$ ), closed symbols ( $\bullet$ ,  $p>0.05$ ).

This work was supported, in part, by the Swiss National Science Foundation grant 3.924-0.87 and the Kanton of Zurich.

- 1 Colucci WS, Wright RF, Braunwald E. New positive inotropic agents in the treatment of congestive heart failure. *N Engl J Med* 1986;314:290-9, 349-58.
- 2 Fisher LA, Kikkawa DO, Rivier JE, et al. Stimulation of noradrenergic sympathetic outflow by calcitonin gene-related peptide. *Nature* 1983;305:534-6.
- 3 Brain SD, Williams TJ, Tippins JR, Morris HR, MacIntyre I. Calcitonin gene-related peptide is a potent vasodilator. *Nature* 1985;313:54-6.
- 4 Gennari C, Fischer JA. Cardiovascular action of calcitonin gene-related peptide in humans. *Calcif Tiss Int* 1985;37:581-4.
- 5 Struthers AD, Brown MJ, Macdonald DWR, et al. Human calcitonin gene-related peptide: a potent endogenous vasodilator in man. *Clin Sci* 1986;70:389-93.
- 6 Holman JJ, Craig RK, Marshall I. Human  $\alpha$ - and  $\beta$ -CGRP and rat  $\alpha$ -CGRP are coronary vasodilators in the rat. *Peptides* 1986;7:231-5.
- 7 McEwan J, Larkin S, Davies G et al. Calcitonin gene-related peptide: a potent dilator of human epicardial coronary arteries. *Circulation* 1986;74:1243-7.
- 8 Franco-Cereceda A, Gennari C, Nami, et al. Cardiovascular effects of calcitonin gene-related peptides I and II in man. *Circ Res* 1987;60:393-7.
- 9 Howden CW, Logue C, Gavin K, Collie L, Rubin PC. Haemodynamic effects of intravenous calcitonin gene-related peptide in man. *Clin Sci* 1988;74:413-8.
- 10 Geppetti P, Maggi CA, Perretti F, Frilli S, Manzini S. Simultaneous release by bradykinin of substance P- and calcitonin gene-related immunoreactivities from capsaicin-sensitive structures in guinea-pig heart. *Br J Pharmacol* 1988;94:288-90.
- 11 Wharton J, Gulbenkian S, Mulder PK, et al. Capsaicin induces a depletion of calcitonin gene-related peptide (CGRP)-immunoreactive nerves in the cardiovascular system of the guinea pig and rat. *J Auton Nerv Syst* 1986;16:289-306.
- 12 Franco-Cereceda A, Lundberg JM, Saria A, Schreimayer W, Triethart HA. Calcitonin gene-related peptide: release by capsaicin and prolongation of the action potential in the guinea-pig heart. *Acta Physiol Scand* 1988;132:181-90.
- 13 Sigris S, Franco-Cereceda A, Muff R, Henke H, Lundberg JM, Fischer JA. Specific receptor and cardiovascular effects of calcitonin gene-related peptide. *Endocrinology* 1986;119:381-9.
- 14 Franco-Cereceda A, Rudehill A, Lundberg JM. Calcitonin gene-related peptide but not substance P mimics capsaicin-induced coronary vasodilation in the pig. *Eur J Pharmacol* 1987;142:235-43.
- 15 Edvinsson L, Fredholm BB, Hamel E, Jansen I, Verrechia C. Perivascular peptides relax cerebral arteries concomitant with stimulation of cyclic adenosine monophosphate accumulation or release of an endothelium-derived relaxing factor in the rat. *Neurosci Lett* 1985;58:213-7.
- 16 Kubota M, Moseley JM, Butera L, Dusting GJ, MacDonald PS, Martin TJ. Calcitonin gene-related peptide stimulates cyclic AMP formation in rat aortic smooth muscle cells. *Biochem Biophys Res Commun* 1985;132:88-94.
- 17 Bruce RA. Evaluation of functional capacity and exercise tolerance of cardiac patients. I. Functional capacity. *Mod Concepts Cardiovasc Dis* 1956;25:321-6.
- 18 Hassan S, Turner P. Systolic time intervals: a review of the method in the non-invasive investigation of cardiac function in health, disease and clinical pharmacology. *Postgrad Med J* 1983;59:423-34.
- 19 Sahn D, DeMaria A, Kisslo J, Weyman A. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* 1978;58:1072-8.
- 20 Smith TW, Butler VP, Haber E. Determination of therapeutic and toxic serum digoxin concentration by radioimmunoassay. *N Engl J Med* 1969;281:1212-6.
- 21 Fischer JA, Blum JW, Binswanger U. Acute parathyroid hormone response to epinephrine in vivo. *J Clin Invest* 1973;52:2434-40.
- 22 Diem K, Lentner C. *Documenta Geigy, wissenschaftliche Tabellen*. 7th ed. Basel: J.R. Geigy 1968:146-97.
- 23 Beglinger C, Born W, Hildebrand P, et al. Calcitonin gene-related peptides I and II and calcitonin: distinct effects on gastric secretion in man. *Gastroenterology* 1988;95:958-65.
- 24 Cachelin AB, de Peyer JE, Kokobun S, Reuter H.  $Ca^{2+}$  channel modulation by 8-bromocyclic AMP in cultured heart cells. *Nature* 1983;304:462-4.
- 25 Tan LB, Murray RG, Littler WA. An analytical method to separate inotropic and vasodilatory drug effects in patients with heart failure. *Cardiovasc Res* 1987;21:625-30.